

## Prevalence and Seasonality of Acute Appendicitis Among Patients Registered at Ibn- Sina General Hospital in Al-Mukalla, Yemen

Abdulla Saleh Alyamani\*

Fauzia Faraj Bamatraf\*\*

### Abstract

Appendicitis is the most common acute surgical condition of the abdomen. Age, sex and seasonal variations have been observed in many studies. The aim of the study is to determine the prevalence and seasonality of acute appendicitis among patients registered in Ibn-Sina general hospital (ISGH) in Al-Mukalla. In this retrospective study. The data was collected from admission records in surgical department of ISGH during the period January 2011 - December 2017. Out of 12580 patients attending surgical ward, 2565 patients with acute appendicitis were collected from records of the hospital. The prevalence of acute appendicitis during the study years was 20.39%. The prevalence of acute appendicitis in males was found to be twice that of females (66.6% and 33.4% respectively). Acute appendicitis was significantly higher among males in the 11-20 age group (70.0%). While in females it was higher in the age group  $\leq 10$  years (51.1%) (p value = 0.000).and decreased with age increase in both males and females. A significant seasonal effect was also observed, with the prevalence of acute appendicitis higher in the winter and autumn (p value = 0.000). In this study obvious differences were observed in the prevalence of acute appendicitis for different seasons, age groups and genders with higher rate in teen age group, male gender and winter and autumn season.

**Keywords:** Acute appendicitis, prevalence, seasonality, Ibn-Sina hospital, Al-Mukalla

### Introduction:

Acute appendicitis is the most common cause of 'acute abdomen. It is defined as a sudden, severe inflammation of the vermiform appendix. It is characterized by abdominal pain, usually localized in the lower right quadrant, with nausea, vomiting and constipation [14]. The etiology of the appendicitis still remains unclear and multifactorial.

The disease has been attributed to a variety of possible causes, including mechanical obstruction [9] inadequate dietary fiber [1] smoking [19], air pollution [10] and familial susceptibility [6,21].

The incidence of appendicitis seems to have risen greatly in the first half of this century, particularly in Europe, America and Australasia, with up to 16% of the population undergoing appendectomy [14]. The rate of acute appendicitis varies among countries. Declining rates of acute appendicitis have been reported in the United States and Europe. In the past 30 years, the incidence has fallen dramatically in these countries, such that the individual lifetime risk of appendectomy is 8.6% and 6.7% among males and females respectively [14]. However, the frequency appears to be increasing in the

developing countries [20,16].

Age, sex and seasonal variations of acute appendicitis have been observed in many studies, but the reasons for these variations are not clear. Acute appendicitis is relatively rare in infants, and is common in childhood and early adult life, reaching the peak of infection in teens and early twenties.. After middle age, the risk of developing appendicitis is quite small. The incidence of appendicitis is equal among males and females before puberty. In teenagers and young adults, the male–female ratio increases to 3:2 at age 25; thereafter, the greater incidence in males declines [14].

Population-based epidemiological studies about the impact of seasonal variables on appendicitis have been published, but so far there is no clear consensus about the connection between the seasonal variations and acute appendicitis [4,15,17,18].

Although acute appendicitis is a common surgical condition, little is known about its epidemiology in Yemen. Therefore, this study was conducted to explore the prevalence, demographic characteristics, and seasonal variation and trends of acute appendicitis among patients attended Ibn Sina general hospital in AL-Mukalla district during the period 1st January 2011 to 31st December 2017.

### Materials and method:

The study was a retrospective descriptive study, relied on data that had been collected through medical records. All patients with Appendicitis

---

\* Surgical Department, College of Medicine and Health Sciences, Hadhrumout University.

\*\* Community Medicine Department, College of Medicine and Health Sciences, Hadhrumout University. Received on 17/3/2018 and Accepted for Publication on 7/5/2018

who registered for treatment at Ibn Sina General Hospital (ISGH) in AL-Mukalla district from 1<sup>st</sup> January 2011 to 31<sup>st</sup> December 2017 were included in this study. The study was carried out at Ibn Sina General Hospital in Al- Mukalla District. ISGH is the main referral hospital that provides health services to the population from the following geographical areas [Hadhrumout, Shabwa and Mahara governorates)

#### Data Analysis:

To ensure the accuracy of data processing, the data were analyzed by using (statistical package for social software SPSS version (19.0), frequencies and percentages for categorical data.

Comparisons were done with the Chi-square test and p-value level <0.05 was considered, significant throughout the study.

#### Results:

During the seven years the total patients attending the surgical wards were 12580; out of them 2565 patients with acute appendicitis were collected from records of the hospital. The prevalence of acute appendicitis during the 7 years was 20.39%. Out of 2565 there were 2055 (80.1%) patients operated with acute appendicitis and 510 (19.9%) were treated conservatively as appear in Table 1.

**Table No (1): Frequency of patients registered at ISGH with acute appendicitis during the years 2011-2017**

Treatment		Frequency	Percent
Operated	Appendectomy	2013	78.5
	Perforation	42	1.6
Conservative	Mass	108	4.2
	Observation	402	15.7
Total		2565	100.0

As shown in Table 2 the distribution of acute appendicitis with respect to gender. Males were found to be twice as high as females (66.6% and 33.4% respectively) of the total number of patients during the seven years of study period. It

was seen also that the number of patients with acute appendicitis is increasing yearly starting from 157 at the year 2011 to 743 at the year 2017.

**Table No (2): Distribution of acute appendicitis related to gender during the study years 2011-2017**

Year	Gender		Total No (%)
	Male No (%)	Female No (%)	
2011	116 (73.9)	41 (26.1)	157 (100.0)
2012	101 (62.7)	60 (37.3)	161 (100.0)
2013	152 (65.5)	80 (34.5)	232 (100.0)
2014	278 (71.1)	113 (28.9)	391 (100.0)
2015	259 (65.7)	135 (34.3)	394 (100.0)
2016	318 (65.3)	169 (34.7)	487 (100.0)
2017	484 (65.1)	259 (34.9)	743 (100.0)
Total	1708 (66.6)	857 (33.4)	2565 (100.0)

Table 3 demonstrates the distribution of acute appendicitis regarding age groups. More than half of recorded patients were at the age group of

11-20 years and the number was decreased when the patients were older. The lowest number was 109 (4.2%) in older than >40 years.

**Table 3: Distribution of acute appendicitis according to age group**

Age group	Frequency	Percent
≤10	223	8.7
11-20	1332	51.9
21-30	729	28.4
31-40	172	6.7
>40	109	4.3
Total	2565	100.0

Using Chi square test, it was found that, there was significant association between patients' age groups and gender in the occurrence of acute appendicitis. The prevalence of acute appendicitis was significantly higher among males in age group 11-20 years 932(70.0%). The

prevalence of acute appendicitis was significantly higher among females in age group ≤ 10 years 114(51.1%). On the other hand the prevalence of acute appendicitis has decreased with increasing of age in both males and females (Table 4).


**Table 4: Association between prevalence of acute appendicitis and age group and gender among study populations**

Age groups (years)	Sex		Total No (%)	p-value
	Male No (%)	Female No (%)		
≤ 10	109 (48.9)	114 (51.1)	223 (100.0)	0.000
11-20	932 (70.0)	400 (30.0)	1332 (100.0)	
21-30	483 (66.3)	246 (33.7)	729 (100.0)	
31-40	113 (65.7)	59 (34.3)	172 (100.0)	
> 40	71 (65.1)	38 (34.9)	109 (100.0)	
Total	1708 (66.6)	857 (33.4)	2565 (100.0)	

\*Chi square  $p < 0.05$  is considered statistically significant

Figure (2) shows the prevalence of acute appendicitis in every month during the study period 2011-2017. It was found that the highest prevalence of registered patients with acute

appendicitis was in cold months; December, November, October and January (260, 256, 248 and 245 cases respectively). The lowest prevalence was in May 156 patients.



As shown in Table 5, there was a significant association between season and prevalence of acute appendicitis. It was clear that the prevalence of acute appendicitis throughout the

most study years was significantly higher in winter and autumn and decreased during spring and summer.

**Table 5: Association between seasons and the prevalence of acute appendicitis**

Year	Season								Total		P. value*
	Winter		Spring		Summer		Autumn		No	%	
	No	%	No	%	No	%	No	%			
2011	39	(24.8)	33	(21.0)	40	(25.5)	45	(28.7)	157	(100.0)	0.000
2012	33	(20.5)	34	(21.1)	54	(33.5)	40	(24.8)	161	(100.0)	
2013	82	(35.3)	48	(20.7)	30	(12.9)	72	(31.0)	232	(100.0)	
2014	97	(24.8)	96	(24.6)	96	(24.6)	102	(26.1)	391	(100.0)	
2015	140	(35.5)	69	(17.5)	100	(25.4)	85	(21.6)	394	(100.0)	
2016	125	(25.7)	110	(22.6)	112	(23.0)	140	(28.7)	487	(100.0)	
2017	176	(23.7)	184	(24.8)	183	(24.6)	200	(26.9)	743	(100.0)	
Total	692	(27.0)	574	(22.4)	615	(24.0)	684	(26.7)	2565	(100.0)	

\*  $p < 0.05$  is considered statistically significant

### Discussion

Despite diagnostic and therapeutic advancement in medicine, appendicitis remains a clinical emergency. In fact, this illness is one of the most common causes of acute abdominal pain. The prevalence of acute appendicitis found in this study (20.39%) is much higher than that found in the study done among the population in the

United States and European countries 7%. [5,12]. On the other hand the prevalence of acute appendicitis among the Yemeni population in the present study is much lower than that in the other developing countries such as India where the prevalence was 30.7% [11]. One of the most striking epidemiological features of appendicitis is the marked variation in

prevalence by age and sex. In this study, the highest prevalence of acute appendicitis was observed among males than females (with a male to female ratio of 2.2: 1), which was consistent with other studies in Canada, Tehran, Nigeria and Ghana where males had higher prevalence of appendicitis more than females [4,15,17,18].

In our study we found a significant association between the prevalence of acute appendicitis and the patients' age. Acute appendicitis was found significantly higher among males in those aged 11-20 years of age group whereas in females the highest occurrence was observed in  $\leq 10$  years of age group. In most of the studies, it is observed that appendicitis is common in the youngest age group which we also observed in our study. These studies such as India (1-20 years age group) United States (10–19 years age group) and Ontario, Canada (10–19 years age group) [11,5,4].

Study conducted by Noudeh et al in Tehran showed that appendicitis is more common in males, in those aged 21-30 years whereas in females the highest occurrence was observed in (11-20 years of age group) [15].

A study conducted by Thomas in Saudi Arabia showed that acute appendicitis was found to be most common between ages 10 to 29 years [23]. While a study conducted by Mungadi et al in Nigeria showed that acute appendicitis is more common in age group 20-29 years [13].

Regarding seasonality our study found a significant association between the occurrence of acute appendicitis and temperature. Seven years period of study (2011-2017) showed seasonal variation of appendicitis with peak in Winter and Autumn and decreased during Spring and Summer. This observation is consistent with the study that was carried out in India which revealed that the highest in Summer and Spring and the lowest in Autumn and Winter [11]

In contrast a seasonal pattern of incidence for acute appendicitis which has been reported by other many studies, which revealed that the highest incidence of acute appendicitis was found in Summer and Spring and the lowest in Autumn and Winter [7,13,23]. The influence of seasonal variables on the incidence of acute appendicitis has been discussed in many studies but yet there is no definite explanation for it [2,3,8,22].

#### **Conclusion**

The prevalence of acute appendicitis during the 7 years was (20.39%). Acute appendicitis was significantly higher among males, in those aged 11-20 years compared to females, in those aged  $\leq 10$  years and has decreased with increasing of age in both males and females. Occurrence of appendicitis was significantly higher during the Winter and Autumn season.

**References:**

- 1- Adamidis D, Roma-Giannikou E, Karamolegou K, Tselalidou E, Constantopoulos A. (2000). Fiber intake and childhood appendicitis. *Int J Food Sci Nutr.* 51: 153-157
- 2- Alder AC, Fomby TB, Woodward WA, Haley RW, Sarosi G, Livingston EH. (2010). Association of viral infection and appendicitis. *Arch Surg.* 145: 63-71
- 3- Alexander P. (2013). Association of monthly frequencies of diverse diseases in the calls to the public emergency service of the city of Buenos Aires during 1999-2004 with meteorological variables and seasons. *Int J Biometeorol.* 57: 83-90
- 4- Al-Omran M, Mamdani M, McLeod RS. (2003). Epidemiologic features of acute appendicitis in Ontario, Canada. *Can J Surg.* 46: 263-268
- 5- Buckius M T., McGrath B, Monk J, Grim R, Theodore B and Ahuja V. (2012). Changing Epidemiology of Acute Appendicitis in the United States: Study Period 1993–2008. *Journal of Surgical Research.* 175: 185–190
- 6- Ergul E. (2007). Heredity and familial tendency of acute appendicitis. *Scand J Surg.* 96: 290-292
- 7- Gallerani M, Boari B, Anania G, Cavallesco G, (2006). Manfredini R. Seasonal variation in onset of acute appendicitis. *Clin Ter.* 157:123-7.
- 8- Ilves I, Fagerström A, Herzig KH, Juvonen P, Miettinen P, Paajanen H. (2014). Seasonal variations of acute appendicitis and nonspecific abdominal pain in Finland. *World J Gastroenterol.* 20(14): 4037-4042
- 9- Jones BA, Demetriades D, Segal I, Burkitt DP. (1985). The prevalence of appendiceal fecaliths in patients with and without appendicitis. A comparative study from Canada and South Africa. *Ann Surg.* 202: 80-82
- 10- Kaplan GG, Dixon E, Panaccione R, Fong A, Chen L, Szyszkowicz M, Wheeler A, MacLean A, Buie WD, Leung T, Heitman SJ, Villeneuve PJ. (2009). Effect of ambient air pollution on the incidence of appendicitis. *CMAJ.* 181: 591-597
- 11- Lohar HP, Calcuttawala MAA, Nirhale DS, Athavale VS, Malhotra M, Priyadarshi N. (2014). Epidemiological aspects of appendicitis in a rural setup, India. *Medical Journal of Patil University.* 7 (6); 753 – 57
- 12- McCahy P. (1994). Continuing fall in the incidence of acute appendicitis. *Ann R Coll Surg Engl.* 76:282-3
- 13- Mungadi IA, Jabo BA, Agwu NP. (2004). A review of appendectomy in Sokoto, North-western Nigeria. *Niger J Med.* 13(3):240e3.
- 14- Norman S W, Bulstrode C J.K and O'Connell P. R O. (2004). *Bailey and Love's Short Practice of Surgery* 25th Edition. 120 pp 4-6.
- 15- Noudeh YJ, Sadigh N and Ahmadnia AY. (2007). Epidemiologic features, seasonal variations and false positive rate of acute appendicitis in Shahr-e-Rey, Tehran. *International Journal of Surgery.* 5: 95-98
- 16- Ofili OP. (1987). Implications of the rising incidence of appendicitis in Africans. *Cent Afr J Med.* 33:243-6.
- 17- Oguntola AS, Adeoti ML, Oyemolade TA. Appendicitis. (2010). Trends in incidence, age, sex, and seasonal variations in South-Western Nigeria. *Ann Afr Med.* 9:213-7.S
- 18- Ohene- Yeboah, M., & Abantanga, FA. (2009). Incidence of acute appendicitis in Kumasi, Ghana. *West Afr J Med.* 28:122-5.
- 19- Oldmeadow C, Wood I, Mengersen K, Visscher PM, Martin NG, Duffy DL. (2008). Investigation of the relationship between smoking and appendicitis in Australian twins. *Ann Epidemiol.* 18: 631-636
- 20- Osman AA. (1974). Epidemiological study of appendicitis in Khartoum. *Int Surg.* 59:218-21.
- 21- Sadr Azodi O, Andrén-Sandberg A, Larsson H. (2009). Genetic and environmental influences on the risk of acute appendicitis in twins. *Br J Surg.* 96: 1336-1340
- 22- Saps M, Blank C, Khan S, Seshadri R, Marshall B, Bass L, Di Lorenzo C. (2008). Seasonal variation in the presentation of abdominal pain. *J Pediatr Gastroenterol Nutr.* 46: 279-284
- 23- Thomas JO. (1998). Acute appendicitis in Buraidah, Saudi Arabia. *Cent Afr J Med.* 44(7):176-8.

## انتشار وموسمية التهاب الزائدة الدودية الحاد بين المرضى المسجلين في مستشفى ابن سينا العام بالمكلا، اليمن

فوزية فرج بامطرف

عبدالله صالح اليماني

### الملخص

لتهاب الزائدة الدودية هو أكثر الحالات الجراحية الحادة شيوعا في البطن. وفي العديد من الدراسات تم ملاحظة العمر والجنس والتغيرات الموسمية. الهدف من الدراسة هو تحديد مدى انتشار وموسمية التهاب الزائدة الدودية الحاد بين المرضى المسجلين في مستشفى ابن سينا العام في المكلا. في هذه الدراسة الاسترجاعية، تم جمع البيانات من سجلات القبول في قسم الجراحة في مستشفى ابن سينا العام بالمكلا خلال الفترة يناير 2011 - ديسمبر 2017. من بين 12580 مريض الذين حضروا جناح الجراحة، تم فرز 2565 مريض من سجلات المرضى والذين يعانون من التهاب الزائدة الدودية الحاد. كانت نسبة التهاب الزائدة الدودية الحاد خلال السبع سنوات 20.39%. ووجد أن انتشار التهاب الزائدة الدودية الحاد في الذكور ضعف الإناث (66.6%) والإناث 33.4%. وكانت التهاب الزائدة الدودية الحاد أعلى بشكل ملحوظ بين الذكور في المجموعة العمرية 11-20 سنة (70.0%). بينما في الإناث كانت أعلى في الفئة العمرية  $\geq 10$  سنوات (51.1%) ( $P=0.000$ )، وانخفضت مع زيادة العمر في كل من الذكور والإناث. ولوحظ أيضا تأثير موسمي كبير، مع انتشار التهاب الزائدة الدودية الحاد أعلى في فصل الشتاء والخريف ( $p = 0.000$ ). في هذه الدراسة لوحظ اختلافات واضحة في التهاب الزائدة الدودية الحاد لمواسم مختلفة، وللفئات العمرية وفي الجنسين مع انتشار أعلى في مجموعة سن المراهقة، وبين الذكور وموسم الشتاء والخريف.

**الكلمات المفتاحية:** التهاب الزائدة الدودية الحاد، موسمية الإصابة، مستشفى ابن سينا، المكلا